

**FINAL STUDY REPORT**PROTOCOL TITLE

Virucidal Efficacy of a Disinfectant for Use on Inanimate Environmental Surfaces

Virus: Herpes simplex virus type 1DATA REQUIREMENTU.S. EPA 40 CFR Part 158,
"Data Requirements for Registration"
Pesticide Assessment Guidelines - Subdivision G, 91-2(f)PRODUCT IDENTITYACCEL TB
Lot 2-3646-REG-US and Lot 3-3647-REG-USPROTOCOL NUMBER

SRC27022304.HSV1

PROJECT NUMBER

A02069

AUTHORMary J. Miller, M.T.
Study DirectorSTUDY COMPLETION DATE

June 30, 2004

PERFORMING LABORATORYATS Labs
1285 Corporate Center Drive, Suite 110
Eagan, MN 55121SPONSORVirox Technologies
6705 Mill Creek Road Unit 4
Mississauga, Ontario L5N5M4SPONSOR REPRESENTATIVEScientific & Regulatory Consultants, Inc.
102 1/2 South Chauncey Street
Columbia City, IN 46725-2306

Page 1 of 14

Project No. A02069
Protocol Number: SRC27022304.HSV1

Virox Technologies
Page 2 of 14




STATEMENT OF NO DATA CONFIDENTIALITY CLAIMS

No claim of confidentiality is made for any information contained in this study on the basis of its falling within the scope of FIFRA Section 10 (d) (1) (A), (B), or (C).

Company: Virox Technologies

Company Agent: Sally Hayes

Agent for Virox Technologies
Title


Signature

Date: 09/20/04

Project No. A02069
Protocol Number: SRC27022304.HSV1

Virox Technologies
Page 3 of 14



GOOD LABORATORY PRACTICE STATEMENT

The study referenced in this report was conducted in compliance with U.S. Environmental Protection Agency Good Laboratory Practice (GLP) regulations set forth in 40 CFR Part 160.

The procedures not performed by or under the direction of ATS Labs are exempt from this Good Laboratory Practice Statement and include: characterization and stability of the compound(s).

Submitter: Sally Hayes
Sally Hayes, Agent for Virox Technologies

Date: 09/20/04

Sponsor: Rhonda Jones
Rhonda Jones, Agent for Virox Technologies

Date: 7-2-04

Study Director: Mary J. Miller
Mary J. Miller, M.T.

Date: 6-30-04

Project No. A02069

Virox Technologies



Protocol Number: SRC27022304.HSV1

Page 4 of 14

QUALITY ASSURANCE UNIT SUMMARY

Study: Virucidal Efficacy of a Disinfectant for Use on Inanimate Environmental Surfaces

The objective of the Quality Assurance Unit is to monitor the conduct and reporting of non-clinical laboratory studies. This study has been performed under Good Laboratory Practice regulations (40 CFR Part 160) and in accordance to standard operating procedures and a standard protocol. The Quality Assurance Unit maintains copies of study protocols and standard operating procedures and has inspected this study on the date(s) listed below. Studies are inspected at time intervals to assure the integrity of the study.

Phase Inspected	Date	Study Director	Management
Critical Phase	April 20, 2004	April 20, 2004	May 3, 2004
Draft Report	April 30, 2004	April 30, 2004	
Final Report	June 28, 2004	June 28, 2004	June 30, 2004

The findings of these inspections have been reported to management and the Study Director.

Quality Assurance Auditor: Rachelle L. Eirson

Date: 06/30/04

Project No. A02069
 Protocol Number: SRC27022304.HSV1

Virox Technologies
 Page 5 of 14



TABLE OF CONTENTS

Title Page	1
Statement of No Data Confidentiality Claims	2
Good Laboratory Practice Statement	3
Quality Assurance Unit Summary	4
Table of Contents	5
Study Personnel	6
General Study Information	7
Test Substance Identity	7
Study Dates	7
Objective	7
Summary of Results	8
Test System	8
Methods	9
Protocol Changes	10
Data Analysis	10
Study Retention	11
References	11
Study Results	12
Study Conclusion	12
Table 1: Virus Control and Test Results	13
Table 2: Cytotoxicity Control Results	13
Table 3: Neutralization Control Results	14

Project No. A02069
Protocol Number: SRC27022304.HSV1

Virox Technologies
Page 6 of 14



STUDY PERSONNEL

STUDY DIRECTOR: Mary J. Miller, M.T.

Professional Personnel Involved:

Douglas G. Anderson, Ph.D.	- President
Karen M. Ramm, B.A.	- Technical Director
Mary J. Miller, M.T.	- Research Scientist II
Katherine A. Paulson, M.L.T.	- Research Assistant II

Project No. A02069
Protocol Number: SRC27022304.HSV1

Virox Technologies
Page 7 of 14



STUDY REPORT

GENERAL STUDY INFORMATION

Study Title: Virucidal Efficacy of a Disinfectant for Use on Inanimate Environmental Surfaces

Project Number: A02069

Protocol Number: SRC27022304.HSV1

Sponsor: Virox Technologies
6705 Mill Creek Road Unit 4
Mississauga, Ontario L5N5M4

Sponsor Representative: Scientific & Regulatory Consultants, Inc.
102 1/2 South Chauncey Street
Columbia City, IN 46725-2306

Testing Facility: ATS Labs
1285 Corporate Center Drive, Suite 110
Eagan, MN 55121

TEST SUBSTANCE IDENTITY

Test Substance: ACCEL TB

Lot/Batch(s): Lot 2-3646-REG-US and Lot 3-3647-REG-US

Test Substance Characterization

Test substance characterization as to content, stability, solubility, storage, etc., is the responsibility of the Sponsor.

STUDY DATES

Date Sample Received: March 11, 2004
Study Initiation Date: April 1, 2004
Experimental Start Date: April 13, 2004
Experimental End Date: April 20, 2004
Study Completion Date: June 30, 2004

OBJECTIVE

The objective of this study was to evaluate the virucidal efficacy of a disinfectant against Herpes simplex virus type 1 according to test criteria and methods approved by the United States Environmental Protection Agency (U.S. EPA) for registration of a product as a virucide.

Project No. A02069
Protocol Number: SRC27022304.HSV1

Virox Technologies
Page 8 of 14



SUMMARY OF RESULTS

Test Substance: ACCEL TB, Lot 2-3646-REG-US and Lot 3-3647-REG-US

Dilution: Ready to use (RTU)

Virus: Herpes simplex virus type 1, ATCC VR-733, Strain F(1)

Exposure Time: One minute

Exposure Temperature: 20±1°C

Organic Soil Load: 5% fetal bovine serum

Efficacy Result: Two lots of ACCEL TB (Lot 2-3646-REG-US and Lot 3-3647-REG-US) met the test criteria specified in the study protocol. The results indicate **complete inactivation** of Herpes simplex virus type 1 under these test conditions as required by the U.S. EPA for claims of virucidal activity.

TEST SYSTEM

- Virus

The F(1) strain of Herpes simplex virus type 1 used for this study was obtained from the American Type Culture Collection, Manassas, VA (ATCC VR-733). The stock virus was prepared by collecting the supernatant culture fluid from infected culture cells. The cells were disrupted and cell debris removed by centrifugation at approximately 2000 RPM for five minutes at approximately 4°C. The supernatant was removed, aliquoted, and the high titer stock virus was stored at ≤ -70°C until the day of use. On the day of use, an aliquot of stock virus (ATS Labs Lot H47) was removed, thawed and refrigerated until use in the assay. The stock virus culture was adjusted to contain 5% fetal bovine serum as the organic soil load. The stock virus tested demonstrated cytopathic effects (CPE) typical of Herpes simplex virus on rabbit kidney cells.
- Test Cell Cultures

Rabbit kidney (RK) cells were obtained from ViroMed Laboratories, inc., Cell Culture Division. Cultures were maintained and used as monolayers in disposable tissue culture labware. On the day of testing, cells were observed as having proper cell integrity and therefore, were acceptable for use in this study.
- Test Medium

Test medium used in this study was Eagle's minimal essential medium (E-MEM) supplemented with 5% heat-inactivated fetal bovine serum (FBS), 10 µg/mL gentamicin, 100 units/mL penicillin, and 2.5 µg/mL amphotericin B (Fungizone).

Project No. A02069
 Protocol Number: SRC27022304.HSV1

Virox Technologies
 Page 9 of 14



The following table lists the test and control groups, the dilutions assayed, and the number of cultures used. See text for a more detailed explanation.

NUMBER OF DILUTIONS AND CULTURES FOR VIRUCIDAL EFFICACY STUDY			
Test or Control Group	Dilutions Assayed (log ₁₀)	Cultures per dilution	Total Cultures
Cell Control	N/A	4	4/group
Dried Virus Control (Group A)	-1,-2,-3,-4,-5,-6,-7,-8	4	32
Sample lot #1 + virus (Group B)	-1,-2,-3,-4,-5,-6,-7,-8	4	32
Sample lot #2 + virus (Group B)	-1,-2,-3,-4,-5,-6,-7,-8	4	32
Cytotoxicity of lot #1 (Group C)	-1,-2,-3,-4,-5,-6,-7,-8	4	32
Cytotoxicity of lot #2 (Group C)	-1,-2,-3,-4,-5,-6,-7,-8	4	32
Non-Virucidal level - lot #1 (Group D)	-1,-2,-3,-4,-5,-6,-7,-8	4	32
Non-Virucidal level - lot #2 (Group D)	-1,-2,-3,-4,-5,-6,-7,-8	4	32

METHODS

- Preparation of Test Substance
 Two lots of ACCEL TB (Lot 2-3646-REG-US and Lot 3-3647-REG-US) were used, undiluted, as received from the Sponsor. The test substance was in solution as determined by visual observation.
- Preparation of Virus Films
 Films of virus were prepared by spreading 0.2 mL of virus inoculum uniformly over the bottoms of three separate 100 x 15mm sterile glass petri dishes. The virus films were dried at 20.0°C in a relative humidity of 54% until visibly dry (20 minutes).
- Sephadex Gel Filtration
 To reduce the cytotoxic level of the virus-disinfectant mixture prior to assay of virus and/or to reduce the virucidal level of the disinfectant, virus was separated from disinfectant by filtration through Sephadex gel. Columns of Sephadex LH-20-100 were equilibrated with phosphate buffered saline containing 1% albumin, centrifuged for three minutes to clear the void volume, loaded with 2.0 mL of virus-disinfectant mixture and immediately passed through the column utilizing the syringe plunger.
- Treatment of Virus Films with Test Substance (GROUP B, TABLE 1)
 For each lot of disinfectant, separate dried virus films were exposed to 2.0 mL of the use dilution for one minute at 20.0°C. Following the exposure time, the plates were scraped with a cell scraper to resuspend the contents of the plate and the virus-disinfectant mixture was immediately passed through a Sephadex column utilizing the syringe plunger in order to detoxify the mixture. The filtrate (10⁻¹ dilution) was then titered by 10-fold serial dilution and assayed for infectivity.

Project No. A02069

Virox Technologies



Protocol Number: SRC27022304.HSV1

Page 10 of 14

5. Treatment of Virus Control Film (GROUP A, TABLE 1)

A virus film was prepared as previously described (paragraph 2). The control film was exposed to 2.0 mL of test medium for the same amount of time as the test film was exposed to the disinfectant. The virus was then scraped and passed through a Sephadex column in the same manner as the test virus and the filtrate (10^{-1} dilution) was then titered by 10-fold serial dilution and assayed for infectivity.

6. Cytotoxicity Assay (GROUP C, TABLE 2)

A 2.0 mL aliquot of the use dilution of each lot of the disinfectant was filtered through a Sephadex column and the filtrate was diluted serially in medium and inoculated into RK cell cultures. Cytotoxicity of the RK cell cultures was scored at the same time as the virus-disinfectant and virus control cultures.

7. Assay of Non-Virucidal Level of Test Substance (GROUP D, TABLE 3)

Each dilution of the Sephadex-filtered disinfectant (disinfectant control for cytotoxicity assay) was mixed with an aliquot of low titer stock virus, and the resulting mixtures of dilutions were assayed for infectivity in order to determine the dilution(s) of disinfectant at which virucidal activity, if any, was retained. Dilutions that showed virucidal activity were not considered in determining the reduction in infectivity by the test substance.

8. Infectivity Assays

The RK cell line, which exhibits CPE in the presence of Herpes simplex virus type 1, was used as the indicator cell line in the infectivity assays. Cells in multiwell culture dishes were inoculated in quadruplicate with 0.1 mL of the dilutions prepared from test and control groups. Uninfected indicator cell cultures (cell controls) were inoculated with test medium alone. Cultures were incubated at 36-38°C in a humidified atmosphere of 5-7% CO₂ in sterile disposable cell culture labware. The cultures were scored periodically for seven days for the absence or presence of CPE, cytotoxicity, and for viability.

9. Statistical Methods: N/A

PROTOCOL CHANGES

Protocol Amendments:

No protocol amendments were required for this study.

Protocol Deviations:

No protocol deviations occurred during this study.

DATA ANALYSIS

Calculation of Titers

Viral and cytotoxicity titers are expressed as $-\log_{10}$ of the 50 percent titration endpoint for infectivity (TCID₅₀) or cytotoxicity (TCD₅₀), respectively, as calculated by the method of Spearman Karber.

$$-\text{Log of 1st dilution inoculated} - \left[\left(\left(\frac{\text{Sum of \% mortality at each dilution}}{100} \right) - 0.5 \right) \times (\text{logarithm of dilution}) \right]$$

Calculation of Log Reduction

Dried Virus Control TCID₅₀ – Test Substance TCID₅₀ = Log Reduction

Project No. A02069
Protocol Number: SRC27022304.HSV1

Virox Technologies
Page 11 of 14



STUDY RETENTION

Record Retention

All of the original raw data developed exclusively for this study shall be archived at ATS Labs, 1285 Corporate Center Drive, Suite 110, Eagan, MN 55121. These original data include, but are not limited to, the following:

1. All handwritten raw data for control and test substances including, but not limited to notebooks, data forms and calculations.
2. Any protocol amendments/deviation notifications.
3. All measured data used in formulating the final report.
4. Memoranda, specifications, and other study specific correspondence relating to interpretation and evaluation of data, other than those documents contained in the final study report.
5. Original signed protocol.
6. Certified copy of final study report.
7. Study-specific SOP deviations made during the study.

Test Substance Retention

The test substance will be discarded following study completion per Sponsor approved protocol. It is the responsibility of the Sponsor to retain a sample of the test material.

REFERENCES

1. Annual Book of ASTM Standards 2000, Section 11 Water and Environmental Technology Volume 11.05 Biological Effects and Environmental Fate: Biotechnology; Pesticides, E1053-97.
2. U.S. Environmental Protection Agency Pesticide Assessment Guidelines, Subdivision G: Product Performance, 91-2(f), November 1982.
3. U.S. Environmental Protection Agency, Registration Division, Office of Pesticide Programs, DIS/TSS-7, November 12, 1981.
4. Diagnostic Procedures for Viral, Rickettsial, and Chlamydial Infections. Schmidt, N.J. and Emmons, R.W. editors. Sixth edition, 1989. p. 18-20.
5. Blackwell, J.H., and J.H.S. Chen. 1970. Effects of various germicidal chemicals on HEP-2 cell culture and Herpes simplex virus. J. AOAC 53:1229-1236.

Project No. A02069

Virox Technologies



Protocol Number: SRC27022304.HSV1

Page 12 of 14

STUDY RESULTS

Results of tests with two lots of ACCEL TB (Lot 2-3646-REG-US and Lot 3-3647-REG-US), ready to use, exposed to Herpes simplex virus type 1, in the presence of a 5% fetal bovine serum soil load at 20.0°C for one minute are shown in Tables 1-3. All cell controls were negative for test virus infectivity. The titer of the dried virus control was 5.25 log₁₀. Following exposure, test virus infectivity was not detected in the virus-test substance mixture for either lot at any dilution tested (≤ 0.5 log₁₀). Test substance cytotoxicity was not observed in either lot at any dilution tested (≤ 0.5 log₁₀). The neutralization control (non-virucidal level of the test substance) indicates that the test substance was neutralized at ≤ 0.5 log₁₀ for both lots. Taking the cytotoxicity and neutralization control results into consideration, the reduction in viral titer was ≥ 4.75 log₁₀ for both lots.

STUDY CONCLUSION

Under the conditions of this investigation, in the presence of a 5% fetal bovine serum soil load, ACCEL TB (Lot 2-3646-REG-US and Lot 3-3647-REG-US), ready to use, demonstrated complete inactivation of Herpes simplex virus type 1 following a one minute exposure time at 20.0°C as required by the U.S. EPA for virucidal label claims.

In the opinion of the Study Director, there were no circumstances that may have adversely affected the quality or integrity of the data.

The use of the ATS Labs name, logo or any other representation of ATS Labs without the written approval of ATS Labs is prohibited. In addition, ATS Labs may not be referred to in any form of promotional materials, press releases, advertising or similar materials (whether by print, broadcast, communication or electronic means) without the express written permission of ATS Labs.

Project No. A02069

Virox Technologies



Protocol Number: SRC27022304.HSV1

Page 13 of 14

TABLE 1: Effects of ACCEL TB (Lot 2-3646-REG-US and Lot 3-3647-REG-US) Following a One Minute Exposure to Herpes Simplex Virus Type 1 Dried on an Inanimate Surface

Dilution	Dried Virus Control (GROUP A)	Herpes simplex virus type 1	
		Lot 2-3646-REG-US (GROUP B)	Lot 3-3647-REG-US (GROUP B)
Cell Control	0 0 0 0	0 0 0 0	0 0 0 0
10 ⁻¹	++++	0 0 0 0	0 0 0 0
10 ⁻²	++++	0 0 0 0	0 0 0 0
10 ⁻³	++++	0 0 0 0	0 0 0 0
10 ⁻⁴	++++	0 0 0 0	0 0 0 0
10 ⁻⁵	++0+	0 0 0 0	0 0 0 0
10 ⁻⁶	0 0 0 0	0 0 0 0	0 0 0 0
10 ⁻⁷	0 0 0 0	0 0 0 0	0 0 0 0
10 ⁻⁸	0 0 0 0	0 0 0 0	0 0 0 0
TCID ₅₀ /0.1 mL	10 ^{5.25}	≤10 ^{0.5}	≤10 ^{0.5}

TABLE 2: Cytotoxicity of ACCEL TB on RK Cell Cultures

Dilution	Cytotoxicity Control	
	Lot 2-3646-REG-US (GROUP C)	Lot 3-3647-REG-US (GROUP C)
Cell Control	0 0 0 0	0 0 0 0
10 ⁻¹	0 0 0 0	0 0 0 0
10 ⁻²	0 0 0 0	0 0 0 0
10 ⁻³	0 0 0 0	0 0 0 0
10 ⁻⁴	0 0 0 0	0 0 0 0
10 ⁻⁵	0 0 0 0	0 0 0 0
10 ⁻⁶	0 0 0 0	0 0 0 0
10 ⁻⁷	0 0 0 0	0 0 0 0
10 ⁻⁸	0 0 0 0	0 0 0 0
TCD ₅₀ /0.1 mL	≤10 ^{0.5}	≤10 ^{0.5}

(+) = Positive for the presence of test virus

(0) = No test virus recovered and/or no cytotoxicity present

Project No. A02069
 Protocol Number: SRC27022304.HSV1

Virox Technologies
 Page 14 of 14



TABLE 3: Non-Virucidal Level of Test Substance (Neutralization Control)

Dilution	Test Virus + Cytotoxicity Control Lot 2-3646-REG-US (GROUP D)	Test Virus + Cytotoxicity Control Lot 3-3647-REG-US (GROUP D)
Cell Control	0 0 0 0	0 0 0 0
10 ⁻¹	++++	++++
10 ⁻²	++++	++++
10 ⁻³	++++	++++
10 ⁻⁴	++++	++++
10 ⁻⁵	++++	++++
10 ⁻⁶	++++	++++
10 ⁻⁷	++++	++++
10 ⁻⁸	++++	++++

(+) = Positive for the presence of test virus after low titer stock virus added (neutralization control)
 (0) = No test virus recovered and/or no cytotoxicity present

Results of the non-virucidal level control indicate that the test substance was neutralized at TCID₅₀ of ≤0.5 log₁₀ for both lots.